

## PAPILIO AGENOR.



### CHARACTER GENERICUS.

*Antennæ* apicem versus crassiores, sæpius clavato-capitatae.

*Alæ* (sedentis) erectæ sursumque conniventes, (volatu diurno.)

*Lin. Syst. Nat.* p. 744.

### CHARACTER SPECIFICUS, &c.

PAPILIO alis dentatis nigris basi sanguineis, primoribus striatis, posticis disco albo maculis nigris.

*Lin. Syst. Nat.* p. 747.

*Clerk. ic. ins. rar.* t. 15.


*Cram. pap.* 3. t. 32. f. A. B.

*Jablonsky pap.* 1. t. 8. f. 3.

---

In variis Asiæ regionibus generatur Papilio Agenor, magnitudine vera in tabula depictus.







5

London, Published Sept 5<sup>th</sup> 1801, by Elizabeth Valden, near St. Newman Street

# AGENOR.



## GENERIC CHARACTER.

*Antennæ* commonly thickening towards the end into a clavated tip.

*Wings* (when at rest) meeting upwards. (Flight diurnal.)

## SPECIFIC CHARACTER.

Black Butterfly with dentated wings red at the base; the lower pair white on the disk, with black spots.

---

This elegant insect is found in several of the Asiatic regions, and is represented on the plate in its natural size.